

Electromagnetic pulses in experiments on high energy and high power laser facilities

F. Consoli¹, V.T. Tikhonchuk^{2,3}, D. Neely^{4,5,6}

1) *ENEA, Fusion and Technologies for Nuclear Safety Department, Frascati, Italy*

E-mail : fabrizio.consoli@enea.it

2) *CELIA, University of Bordeaux, CNRS, CEA, 33405 Talence, France*

3) *ELI-Beamlines, Institute of Physics, Czech Academy of Sciences, Czech Republic*

4) *Central Laser Facility, Rutherford Appleton Lab.y, Chilton, Didcot, STFC, UKRI, UK*

5) *AWE plc, Aldermaston, Reading, Berkshire RG7 4PR, UK*

6) *Department of Physics, Scottish Univ. Physics Alliance, Univ. of Strathclyde, Glasgow, UK*

The generation of electromagnetic fields of remarkable intensity in the radiofrequency-microwave range is commonly observed in experiments where high energy and high power lasers interact with matter. These transient electromagnetic pulses (EMPs) depend on laser energy and intensity [1]. They can carry a significant portion of the incoming laser energy and be so powerful to represent a potential serious danger for any electronic device placed inside or even outside the experimental vacuum chamber. For this reason, understanding the origin and the mitigation of these electromagnetic fields is of primary importance for the operation of existing laser facilities for inertial confinement fusion and laser-plasma acceleration. Of course, EMPs represent an even more severe issue for future lasers installations with higher energy and power. On the other hand, they can be positively used for a large number of applications.

Several mechanisms are now recognized as sources of these EMP fields [2]. Target positive charging caused by fast-electron emission due to the laser-plasma interaction is the main one. This fast charging induces high neutralization currents from the conductive walls of the vacuum chamber [3]. The complex picture of the field distribution within the experimental chamber strongly depends on the physical localization and on the characteristics of each of the source processes, but is also affected by the expanding plasma and particle beams emitted from the target [1,2]. An up-to-date presentation of the problems related to generation, detection and mitigation of the strong electromagnetic pulses created in the interaction of high-power, high-energy laser pulses with different types of solid targets will be here given. This will include the activities on the EMP topic performed by the main international research centers active in this field.*

References

[1] F. Consoli et al, High Power Laser Science and Engineering 8 (2020).

[2] F. Consoli, Philosophical Transactions of the Royal Society A 379, 20200022 (2020)

[3] J.-L. Dubois et al, Physical Review E 89, 013102 (2014).

*This work has been carried out within the framework of the EUROfusion Consortium and funded from the Euratom research and training programme 2014-2018 and 2019-2020 under grant agreement No. 633053.